arXiv:hep-ph/9604258v1 7 Apr 1996

THE STRONG $g_{B^{**}B\pi}$ COUPLING CONSTANT IN FULL QCD.

T.M.ALIEV [*], and M. SAVCI
Physics Department, Middle East Technical University
06531 Ankara, Turkey

February 1, 2008

Abstract

To leading order in α_s, the leading and non-leading $1/m_b$ corrections to the excited B^{**} meson coupling $g_{B^{**}B\pi}$ is calculated in the framework of QCD spectral moment sum rules in the full theory. Our prediction is in good agreement with the $light-cone$ QCD sum rule result.

---

[*] Permanent address: Institute of Physics, Azerbaijanian Academy of Sciences, Baku, Azerbaijan

1 Introduction

At present the physics of hadrons containing heavy quarks lies in the focus of experimentalists and theoretists. From experimental point of view it is connected in the first hand, to the the observed exclusive $B \to K^{*}\gamma$ [1] and inclusive decays $B \to X_S\gamma$ [2], which clearly demonstrated the existence of $FCNC$ and in turn, opens a window for the determination of the *Cabibbo-Kobayashi-Maskawa* (CKM) matrix elements V_{tb} and V_{ts}. From theoretical point of view the reason is that the investigation of these decays allows one to check the predictions of CKM at loop level. At the same time these decays form the basis of a very rich *"laboratory framework"* for theoretical researches. For example, in the infinite heavy quark mass limit ($m_Q \to \infty$), QCD reveals symmetries that are not present in the finite mass theory: chiral $SU_L(3) \otimes SU_R(3)$ symmetry, spin-flavour symmetry [3]. These symmetries allow the derivation of a simple relation among the formfactors of the different processes (see for review [4]) and build an effective heavy-light Lagrangian. This effective Lagrangian widely applied in the investigations of B and D meson physics. This approach allows us to include the positive parity (Qq) meson to the effective Lagrangian according to the value of the value of the angular momentum of the ligth degree of freedom ($s_l^P = \frac{1^+}{2}, \frac{3^+}{2}$). The heavy quark effective theory [5, 6] predicts the existence of two multiplets, the first one containing 0^+ and 1^+ and the second 1^+ , 2^+ states mesons.

In [7] the strong coupling constant $g_{B^{**}B\pi}$, where B^{**} is the 0^+ p-wave in the framework of the *classical* and *light cone* sum rules, is investigated. Predictions of the both approaches differ one another by a factor ~ 1.5. This article is devoted to the study of the large $(1/m_b)$ behavior of the $g_{B^{**}B\pi}$ coupling constants and estimate their values in the double momentum version of the QCD sum rules. Contrary to the double exponential version of sum rules, this version is advantageous in the analysis of the three point function, as it prevents the blow up of the QCD series, when m_Q is large. Note that the coupling constant $g_{B^{*}B\pi}$is calculated in the framework of the above mentioned method in [8].

2 Calculation of of the excited $g_{B^{**}B\pi}$ coupling

We start our calculation by considering the following correlator:

$$V = -\int d^4x \; d^4y \; e^{i(p'y - px)} \langle 0 | \mathcal{T} \left\{ J_{B^{**}}(x) J_5(0) J_B(y) \right\} | 0 \rangle \;, \tag{1}$$

where the interpolating currents

$$\begin{aligned} J_{B^{**}} &= m_b \bar{u} b \ , \\ J_5 &= (m_u + m_d) \bar{u} \gamma_5 d \qquad\qquad and \\ J_B &= (m_d + m_b) \bar{d} \gamma_5 b \ , \end{aligned} \tag{2}$$

are described 0^+ excited B meson state in the $s_l^P = \frac{1^+}{2}$ doublet, π-meson and B-meson states. Acoording to QCD ideology this correlator is calculated theoretically in the deep-Eucledian region and in terms of physical states. In order perform this calculation we remind that the correlator obeys the double dispersion relation

$$V(p, p', q) = -\frac{1}{4\pi^2} \int_{m_b^2}^{\infty} \frac{ds}{s - p^2} \int_{m_b^2}^{\infty} \frac{ds'}{s' - p'^2} \, \mathrm{Im} V(s, s') \quad + \quad ... \tag{3}$$

Since m_b is much larger than the QCD scale Λ, it is natural to use the momentum version of sum rules. Acting on both sides of Eq. (3) by the double momentum operator, $\left(-\frac{d}{dp^2}\right)^n \left(-\frac{d}{dp'^2}\right)^n$ over variables p^2, p'^2, and after differentiation setting $p^2 = p'^2 = 0$, we get

$$\mathcal{M}^{(n,n')} = -\frac{1}{4\pi^2} \int_{m_b^2}^{\infty} \frac{ds}{s^{n+1}} \int_{m_b^2}^{\infty} \frac{ds'}{s'^{n'+1}} \, \mathrm{Im} V(s, s') \ . \tag{4}$$

The perturbative part of the spectral function can be calculated from the Feynman diagrams by the help of the Gutkovsky rule, i.e., replacing propagators

$$\frac{1}{(p_i^2 - m_i^2)} \rightarrow -2\pi i \delta(p_i^2 - m_i^2) \ . \tag{5}$$

After carrying standard calculations for spectral density we get,

$$-\frac{1}{4\pi^2} \mathrm{Im} V(s, s') = (m_u + m_d) m_b^3 \, \frac{N_c}{8\pi^2} \, \frac{Q^2}{\lambda^{1/2}(s, s', Q^2)} \ , \tag{6}$$

where $Q^2 \equiv -q^2 \geq 0$ is the pion momentum squared and

$$\lambda = (s + s' + Q^2)^2 - 4ss' \ . \tag{7}$$

Integration region defined by the following inequality:

$$(s - m_b^2)(s' - m_b^2) \geq Q^2 m_b^2. \tag{8}$$

The physical part of the sum rules is parametrized using the usual duality anzats: lowest resonances + QCD continuum, which usually modelled as bare loop, starting from the thresholds s_0 and s_0'. Transferring the the continuum contribution to the theoretical part we obtain the following QCD sum rules

$$g_{B^{**}B\pi}\ \frac{\sqrt{2}m_{B^{**}}^2 f_{B^{**}}}{m_{B^{**}}^{2(n+1)}}\ \frac{\sqrt{2}m_B^2 f_B}{m_B^{2(n'+1)}}\ \frac{\sqrt{2}m_\pi^2 f_\pi}{m_\pi^2+Q^2} \simeq -\frac{1}{4\pi^2}\int_{m_b^2}^{s_0}\frac{ds}{s^{n+1}}\int_{m_b^2}^{s_0'}\frac{ds'}{s'^{n'+1}}\ \mathrm{Im}V(s,s'), \tag{9}$$

In deriving the left hand side of the sum rules we use the following definitions:

$$\begin{aligned}
g_{B^{**}B\pi} &= \langle B(p')\pi(q)|B^{**}(p)\rangle \\
\langle 0|J_5|\pi\rangle &= \sqrt{2}f_\pi m_\pi^2 \\
\langle 0|J_B|B\rangle &= \sqrt{2}f_B m_B^2 \\
\langle 0|J_{B^{**}}|B^{**}\rangle &= \sqrt{2}f_{B^{**}}m_{B^{**}}^2
\end{aligned} \tag{10}$$

In the $m_b \to \infty$ case it is convenient to work with the non-relativistic variables E and E' defined as

$$s=(E+m_b)^2 \qquad \text{and} \qquad s'=(E'+m_b)^2, \tag{11}$$

and introduce the new variables:

$$x=E-E' \qquad \text{and} \qquad y=\frac{1}{2}(E+E')\ . \tag{12}$$

The integration region over new variables x and y is defined from Eq. (8). In further analysis we shall set $n=n'=n_3$. By keeping the non-leading terms in the expansion to the leading order in α_s:

$$\begin{aligned}
\mathcal{M}^{(n,n')} \simeq\ & (m_u+m_d)\ \frac{N_c}{m_b^{4n_3}}\ \frac{Q^2}{4\pi^2}\left\{\int_{-E_c^{B^{**}}}^{0}dx\int_{\frac{1}{2}\sqrt{x^2+Q^2}}^{E_c^{B^{**}}+\frac{x}{2}}dy+\int_0^{E_c^B}dx\int_{\frac{1}{2}\sqrt{x^2+Q^2}}^{E_c^B-\frac{x}{2}}dy\right\} \\
& \left[\frac{1}{(x^2+Q^2)^{1/2}}\Big[1-(4n_3+3)\frac{y}{m_b}+2(n_3+1)(4n_3+3)\frac{y^2}{m_b^2}+\right. \\
& \left.(2n_3+1)\frac{x^2}{4m_b^2}-\frac{Q^2}{8m_b^2}\Big]\right]\ .
\end{aligned} \tag{13}$$

Following [7, 8] , we shall use in our analysis the lowest order expression in α_s for the decay constants $f_{B^{**}}$ and f_B from the 2-point momentum sum rule,

$$\begin{aligned}
f_B^2 &\simeq \frac{\left(E_c^B\right)^3}{2\pi^2}\frac{1}{m_B}\left(\frac{m_B}{m_b}\right)^{2n_2-1}\left\{1-\frac{3}{2}(n_2+1)\frac{E_c^B}{m_b}-\frac{\pi^2}{2}\ \frac{\langle\bar{d}d\rangle}{(E_c^B)^3}\right\}\ , \\
f_{B^{**}}^2 &\simeq \frac{\left(E_c^{B^{**}}\right)^3}{2\pi^2}\frac{1}{m_B^{**}}\left(\frac{m_B^{**}}{m_b}\right)^{2n_2-1}\left\{1-\frac{3}{2}(n_2+1)\frac{E_c^{B^{**}}}{m_b}+\frac{\pi^2}{2}\ \frac{\langle\bar{d}d\rangle}{(E_c^{B^{**}})^3}\right\}\ .
\end{aligned} \tag{14}$$

From the numerical analysis of f_B including the α_s correction one can obtain for the effective value of E_c^B [8]:

$$E_c^B \simeq (1.3 \pm 0.1) \text{ GeV} . \tag{15}$$

Performing similiar calculation using Eq. (14) and the results cited in [9] for the effective value $E_c^{B^{**}}$ of $f_{B^{**}}$ we get,

$$E_c^{B^{**}} \simeq (1.8 \pm 0.1) \text{ GeV} . \tag{16}$$

Using the recently reported experimental results [10, 11] $m_{B^{**}} = (5.732 \pm 0.002)$ MeV and $m_B = 5.279$ MeV we conclude that $m_{B^{**}} - m_B = \Delta \simeq 0.5$ GeV. From comparision of Eqs. (15) and (16) we have,

$$E_c^{B^{**}} \simeq E_c^B + \Delta . \tag{17}$$

Taking into consideration these values of $E_c^{B^{**}}$ and m_B^{**} and following [12], we minimize the n-dependence of the results, by requiring that the leading term is n-independent. This leads us to the following condition:

$$4n_3 = 2n_2 - 1 . \tag{18}$$

From analysis of two point function it is found that $n_2 \simeq 4 - 5$ (see [13]) . Evaluating different integrals, the results can be written in the following good approximation,

$$g_{B^{**}B\pi} \simeq g_{B^{**}B\pi}^{LO} \left\{ 1 + \frac{E_c^B}{m_b} \right\} , \tag{19}$$

where:

$$g_{B^{**}B\pi}^{LO} \simeq \frac{N_c(m_u + m_d)m_B^2}{4\sqrt{2} f_\pi m_\pi^2} \; \mathcal{I}_0 \tag{20}$$

and

$$\mathcal{I}_0 = \frac{Q^4}{m_B \left(E_c^B\right)^3} \left[\int_0^{E_c^{B^{**}}} dx \int_{\frac{1}{2}\sqrt{x^2+Q^2}}^{E_c^{B^{**}} - \frac{x}{2}} + \int_0^{E_c^B} dx \int_{\frac{1}{2}\sqrt{x^2+Q^2}}^{E_c^B - \frac{x}{2}} \right] \frac{dy}{\left(x^2 + Q^2\right)^{1/2}} . \tag{21}$$

In the region $1 \leq Q^2 \leq 4$ this integral has a maximum at $Q^2 \simeq 2.5$ GeV^2. Using this value of Q^2, numerical integration of $\mathcal{I}_0$ for different values of E_c^B can be parametrized by following interpolating formula

$$\mathcal{I}_0 = (0.175 \pm 0.001) E_c . \tag{22}$$

Using

$$E_c^B = E_c^\infty \left(1 - \frac{E_c^B}{2m_b} \right), \quad \text{(see for example [8])} , \tag{23}$$

we finally get

$$g_{B^{**}B\pi} \simeq \frac{3m_B^2}{4\sqrt{2}f_\pi} g^\infty \left\{ 1 + \frac{E_c^B}{2m_b} \right\} \ . \tag{24}$$

Using the result $g^\infty = (0.15 \pm 0.03)$, that is given in [8] and taking into consideration the $\frac{1}{m_b}$ correction for the *physical* $B - meson$ (which is of the order of ±28% at the b-mass), we get,

$$g_{B^{**}B\pi} = 24 \pm 7 \ \ GeV \ , \tag{25}$$

where the error in the above result takes into account the effect of radiative corrections to f_B. As mentioned in Introduction, the value of the strong coupling constant $g_{B^{**}B\pi}$ that is calculated in [7] within the framework of the *classical* and *light-cone sum rule* techniques are ,

$$\begin{aligned} g_{B^{**}B\pi} &= 13.3 \pm 4.8 \ \ GeV, \\ g_{B^{**}B\pi} &= 21 \pm 7 \ \ GeV, \end{aligned} \tag{26}$$

respectively. From comparision of Eqs. (25) and (26) we see that our prediction on $g_{B^{**}B\pi}$ is in good agrrement with the *light-cone* QCD sum rule prediction on the same quantity.

References

[1] R. Ammar *et. al.*, (**CLEO** Colloboration), *Phys. Rev. Lett.* **71** (1994) 674.

[2] M. S. Alam *et. al.*, (**CLEO** Colloboration), *Prep.* CLNS 94-1314 (1994).

[3] N. Isgur and M. B. Wise, *Phys. Lett.* **B232** (1989) 113; **237** (1990) 527.
M. B. Voloshin and M. A. Shifman, *Sov. J. Nucl. Phys.* **45** (1987) 292; **47** (1988) 511.
H. D. Politzer and M. B. Wise, *Phys. Lett.* **B206** (1988) 681; **B208** (1988) 504.
B. Grinstein, *Nucl. Phys.* **B339** (1990) 253.
A. F. Falk, H. Georgi, B. Grinstein and M. B. Wise *ibid* **B343** (1990) 1.

[4] M. Neubert, *Phys. Reports* (1994).

[5] N. Isgur and M. B. Wise, *Phys. Rev. Lett.* **66** (1991) 1130; *Phys. Rev.* **D43** (1991) 819.

[6] A. F. Falk and M. Luke, *Phys. Lett.* **B292** (1992) 119; U. Kilian, J. Körner and D. Pirjol, *ibid* **B288** (1992) 360.

[7] P. Colangelo, F. De Fazio and G. Nardulli, *Phys. Rev.* **D52** (1995) 6422.

[8] H. G. Dosch and S. Narison, *Phys. Lett.* **B368** (1996) 163.

[9] P. Colangelo, G. Nardulli, A. A. Orchninnikov and N. Paver, *Phys. Lett.* **B269** (1991) 201.

[10] **DELPHI** Colloboration, R. Abreu *et. al.*, *Phys. Lett.* **B345** (1995) 598.

[11] **OPAL** Colloboration, R. Akers *et. al.*, *Z. Phys.* **C66** (1995) 19.

[12] S. Narison, *Phys. Lett.* **B345** (1995) 166; *Phys. Lett.* **B337** (1994) 163; *Phys. Lett.* **B327** (1994) 354.

[13] S. Narison, *Phys. Lett.* **B198** (1987) 104; *Phys. Lett.* **B218** (1989) 238; *Phys. Lett.* **B308** (1993) 308.